

Time: 10 minutes

1. Find the product of the real solutions of $x^{2010} = 1$.
 2. In a cube whose edges have length $10\sqrt{6}$, $ABCD$ and $EFGH$ are opposite faces. Find the distance from A to the center of $EFGH$.
-

Time: 10 minutes

3. Find the least positive integer n such that $1040n$ is a square.
 4. The lengths of the medians of a triangle are 9, 9, and 12. The area of the triangle can be expressed as $p\sqrt{q}$, where p and q are positive integers and q is not divisible by the square of a prime. Find $p + q$.
-

Time: 10 minutes

5. The coordinates of point A are $(12, 77)$ and the coordinates of point B are $(68, -53)$. Point P is on \overline{AB} so that $AP:PB = 3:7$. The coordinates of P are (m, n) . Find $10(m + n)$.
6. Find the coefficient of x^5 in the expansion of $(1 + x + x^2)^5$.

Solutions for Contest #2

1. The two real solutions are 1 and -1 , and their product is -1 .
2. Let O be the center of $EFGH$. Without loss of generality, let $ABFE$ be a face of the cube, and let P be the midpoint of \overline{EF} . Draw \overline{OP} and \overline{PA} . Then
$$PA^2 = PE^2 + EA^2 = (10\sqrt{6})^2 + (5\sqrt{6})^2 = 750, \text{ and so}$$
$$AO^2 = AP^2 + PO^2 = 750 + (5\sqrt{6})^2 = 900. \text{ Thus } AO = 30.$$
3. Notice that $1040 = 2^4 \cdot 5 \cdot 13$. The least possible value of n is therefore $5 \cdot 13 = 65$.
4. Label the triangle ABC , let K be its area, let \overline{AD} and \overline{BE} be the medians of length 9, and let \overline{CF} be the median (and altitude) of length 12. Let G be the centroid of the triangle (the point of intersection of the medians). Because the medians of a triangle divide each other in the ratio 2:1, $AG = 6$ and $GF = 4$. Use the Pythagorean Theorem to conclude that $AF = 2\sqrt{5}$. Then the area of right triangle AFG is $(1/2)2\sqrt{5} \cdot 4 = 4\sqrt{5}$, and so $K = 6 \cdot 4\sqrt{5} = 24\sqrt{5}$. Thus $p + q = 29$.
5. Each of the coordinates of P is the weighted average of the corresponding coordinates of A and B . In particular, the coordinates of P are
$$\left(\frac{7 \cdot 12 + 3 \cdot 68}{10}, \frac{7 \cdot 77 + 3 \cdot -53}{10} \right) = (28.8, 38). \text{ Thus } 10(m + n) = 668.$$
6. The given product contains five factors of $(x^2 + x + 1)$. To expand the product, you must choose one term from 1, x and x^2 in each of the five factors. If you choose a 1's, b x 's and c x^2 's, then $a + b + c = 5$. In order for the product of the terms to be x^5 , you must have $1^a \cdot x^b \cdot (x^2)^c = x^5$, that is, $b + 2c = 5$. Thus $a = c$, and so you must choose either no 1's, five x 's and no x^2 's; one 1, three x 's and one x^2 ; or two 1's, one x and two x^2 's.

Count the three cases separately. In the first case, there is one way to choose no 1's, five x 's and no x^2 's. In the second case, there are 5 ways to choose a 1, then 4 ways to choose an x^2 for a total of 20 ways to choose one 1, three x 's and one x^2 . In the third case, there are 5 ways to choose one x , then $\binom{4}{2} = 6$ ways to choose two 1's for a total of 30 ways to choose two 1's, one x and two x^2 's. Thus the coefficient of x^5 is $1 + 20 + 30 = 51$.

Alternatively, we can count as follows:

Case I: choosing one x from each of the five trinomial factors, only one way

Case II: Choosing three x 's, one 2 , and one 1 is analogous to counting the number of arrangements of the letters in the word GEESE. $5!/(1!3!1!)$

Case III: Choosing two 2 , one x and two 1 's is analogous to counting the number of arrangements of the letters in the word MAMAS. $5!/(2!1!2!)$